## Midterm COMP 2804

## February 27, 2014

- All questions must be answered on the scantron sheet.
- Write your name and student number on the scantron sheet.
- You do not have to hand in this examination paper.
- This is a closed-book exam.
- Calculators are not allowed.

Marking scheme: Each of the 17 questions is worth 1 mark.

• Newton:  $(x+y)^n = \sum_{k=0}^n \binom{n}{k} x^{n-k} y^k$ .

- 1. On a table, you see three types of fruit: apples, bananas, and oranges. There are  $m \ge 2$  apples,  $n \ge 2$  bananas, and  $k \ge 2$  oranges. How many ways are there to choose 7 pieces of fruit, if you must take at least two pieces of each type?
  - (a)  $\binom{m+n+k}{7} (m+n+k)$
  - (b)  $\binom{m+n+k}{7} \binom{m}{2} \binom{n}{2} \binom{k}{2}$
  - (c)  $\binom{m}{3}\binom{n}{2}\binom{k}{2} + \binom{m}{2}\binom{n}{3}\binom{k}{2} + \binom{m}{2}\binom{n}{2}\binom{k}{3}$
  - (d)  $\binom{m}{2} \binom{n}{2} \binom{k}{2} (m+n+k)$
- 2. Consider 9 boys and 15 girls. How many ways are there to arrange these 24 people on a line if all boys stand next to each other and all girls stand next to each other?
  - (a)  $\frac{24!}{9!15!}$
  - (b)  $\binom{24}{9}(9!)(15!)$
  - (c) (9!)(15!)
  - (d) 2(9!)(15!)
- 3. Let S be a set of size 37, and let x, y, and z be three distinct elements of S. How many subsets of S are there that contain x and y, but do not contain z?
  - (a)  $2^{33}$
  - (b)  $2^{34}$
  - (c)  $2^{35}$
  - (d)  $2^{37} 2^{35} 2^{36}$
- 4. Let S be a set of size 37, and let x, y, and z be three distinct elements of S. How many subsets of S are there that contain x or y, but do not contain z?
  - (a)  $2^{36} 2^{34}$
  - (b)  $2^{36} 2^{35}$
  - (c)  $2^{37} 2^{34}$
  - (d)  $2^{37} 2^{35}$

5. A password consists of 12 or 13 characters, each character being one of the 10 digits  $0, 1, 2, \ldots, 9$ . A password must contain the digit 7 at least once. How many passwords are there?

(a) 
$$10^{12} + 10^{13} - 9^{12} - 9^{13}$$

(b) 
$$12^{10} + 13^{10} - 12^9 - 13^9$$

(c) 
$$10^{12} + 10^{13} - 7^{12} - 7^{13}$$

(d) 
$$12^{10} + 13^{10} - 12^7 - 13^7$$

6. Let  $n \ge 7$  and  $k \ge 1$  be integers, let A be the set of all bitstrings of length n that contain exactly seven 0s, and let B be the set of all bitstrings of length k that contain at least one 1. Assume there exists a one-to-one function  $f: A \to B$ . Which of the following is true?

(a) 
$$2^k - 1 < \binom{n}{7}$$

(b) 
$$2^k - 1 \ge \binom{n}{7}$$

(c) 
$$2^k - 1 < 2^n / \binom{n}{n-7}$$

(d) 
$$2^k - 1 \ge 2^n / \binom{n}{n-7}$$

- 7. What is the coefficient of  $x^9y^{16}$  in the expansion of  $(7x + 21y)^{25}$ ?
  - (a)  $\binom{25}{16} 7^{16} 21^9$
  - (b)  $\binom{16}{25}7^921^{16}$
  - (c)  $\binom{25}{16} 7^{25} 3^{16}$
  - (d) none of the above
- 8. How many solutions are there to the equation  $x_1 + x_2 + x_3 = 17$ , where  $x_1 \ge 0$ ,  $x_2 \ge 0$ , and  $x_3 \ge 0$  are integers?
  - (a)  $\binom{19}{16}$
  - (b)  $\binom{19}{17}$
  - (c)  $\binom{20}{16}$
  - (d)  $\binom{20}{17}$

9. How many strings can be obtained by rearranging the letters of the word

## POOPERSCOOPER

- (a) 13!
- (b)  $\binom{13}{4} \binom{9}{2} \binom{7}{2} \binom{5}{3}$
- (c)  $\binom{13}{4} \binom{9}{3} \binom{6}{2} \binom{4}{2}$
- (d)  $\binom{13}{1}\binom{12}{4}\binom{8}{2}\binom{6}{1}\binom{5}{2}\binom{3}{3}$
- 10. The function  $f: \mathbb{N} \to \mathbb{N}$  is defined by

$$f(0) = 2$$
  
 $f(n+1) = f(n) + 6n - 2 \text{ for } n \ge 0$ 

What is f(n)?

- (a)  $f(n) = 3n^2 5n + 2$
- (b)  $f(n) = 3n^2 + 5n + 2$
- (c)  $f(n) = 2n^2 5n + 2$
- (d)  $f(n) = 2n^2 + 5n + 2$
- 11. Consider the following recursive algorithm Fib, which takes as input an integer  $n \geq 0$ :

**Algorithm** 
$$Fig(n)$$
:

if 
$$n = 0$$
 or  $n = 1$ 

then 
$$f = n$$

else 
$$f = Fib(n-1) + Fib(n-2)$$

endif;

return f

When running Fib(7), how many calls are there to Fib(3)?

- (a) 4
- (b) 5
- (c) 6
- (d) 7

12. The Fibonacci numbers are defined as follows:  $f_0 = 0$ ,  $f_1 = 1$ , and  $f_n = f_{n-1} + f_{n-2}$  for  $n \ge 2$ .

Consider again the recursive algorithm Fib, which takes as input an integer  $n \geq 0$ :

```
Algorithm FIB(n):

if n = 0 or n = 1

then f = n

else f = FIB(n - 1) + FIB(n - 2)

endif;

return f
```

For  $n \geq 3$ , run algorithm Fib(n) and let  $a_n$  be the number of times that Fib(2) is called. Which of the following is true?

- (a) For  $n \ge 3$ ,  $a_n = f_{n-1}$
- (b) For  $n \geq 3$ ,  $a_n = f_n$
- (c) For  $n \ge 3$ ,  $a_n = f_{n+1}$
- (d) For  $n \ge 3$ ,  $a_n = -1 + f_n$
- 13. Let  $B_n$  be the number of bitstrings of length n that do not contain 111. Which of the following is true?
  - (a)  $B_n = B_{n-1} + B_{n-2} + 2^{n-3}$
  - (b)  $B_n = B_{n-1} + B_{n-2} + 2^{n-3} B_{n-3}$
  - (c)  $B_n = B_{n-1} + B_{n-2} + B_{n-3}$
  - (d)  $B_n = B_{n-1} + B_{n-2} + B_{n-3} + 2^{n-4}$
- 14. A standard deck of 52 cards has 4 Kings. Consider a hand of 9 cards, chosen uniformly at random. What is the probability that there are exactly two Kings in this hand?
  - (a)  $1 {\binom{48}{7}}/{\binom{52}{9}}$
  - (b)  $\left\{ \binom{4}{2} + \binom{48}{7} \right\} / \binom{52}{9}$
  - (c)  $\binom{52}{9} / {\binom{4}{2} \binom{48}{7}}$
  - (d)  $\binom{4}{2}\binom{48}{7}/\binom{52}{9}$

15. We choose a bitstring of length 25 uniformly at random. What is the probability that this string contains at least two 1s?

(a) 
$$1 - (1/2)^{25} - 25(1/2)^{25}$$

(b) 
$$1 + (1/2)^{25} - 25(1/2)^{25}$$

(c) 
$$\sum_{k=2}^{25} {25 \choose k} (1/2)^k$$

- (d) none of the above
- 16. Consider three people, each one having a uniformly random birthday (out of 365 days; we ignore leap years). What is the probability that at least two of them have the same birthday?

(a) 
$$1 - \frac{365^2}{364 \cdot 363}$$

(b) 
$$1 - \frac{364 \cdot 363}{365^2}$$

(c) 
$$1 - {3 \choose 2}/365^3$$

(d) 
$$1 - \left\{ \binom{3}{2} + \binom{3}{3} \right\} / 365^3$$

- 17. What is Simon Pratt's favorite drink?
  - (a) Herbal tea
  - (b) India Pale Ale
  - (c) Poutine
  - (d) None of the above, because Simon doesn't like beer