

# COMP 3804 — Assignment 4

**Due:** Thursday April 6, 23:59.

## Assignment Policy:

- Your assignment must be submitted as one single PDF file through Brightspace.


Use the following format to name your file:

LastName\_StudentId\_a4.pdf

- **Late assignments will not be accepted. I will not reply to emails of the type “my internet connection broke down at 23:57” or “my scanner stopped working at 23:58”, or “my dog ate my laptop charger”.**
- You are encouraged to collaborate on assignments, but at the level of discussion only. When writing your solutions, you must do so in your own words.
- Past experience has shown conclusively that those who do not put adequate effort into the assignments do not learn the material and have a probability near 1 of doing poorly on the exams.
- When writing your solutions, you must follow the guidelines below.
  - You must justify your answers.
  - The answers should be concise, clear and neat.
  - When presenting proofs, every step should be justified.

**Question 1:** Write your name and student number.

**Question 2:** Let  $K \geq 3$  be an integer. A  $K$ -kite is a graph consisting of a clique of size  $K$  and a path with  $K$  vertices that is connected to one vertex of the clique; thus, the number of vertices is equal to  $2K$ . In the figure below, the graph with the black edges forms a 5-kite.



The *kite problem* is defined as follows:

$$\text{KITE} = \{(G, K) : \text{graph } G \text{ contains a } K\text{-kite}\}.$$

Prove that the language KITE is in **NP**.

**Question 3:** The *clique problem* is defined as follows:

$$\text{CLIQUE} = \{(G, K) : \text{graph } G \text{ contains a clique of size } K\}.$$

Prove that  $\text{CLIQUE} \leq_P \text{KITE}$ , i.e., in polynomial time, CLIQUE can be reduced to KITE.

**Question 4:** The *subset sum problem* is defined as follows:

$$\text{SUBSETSUM} = \{(S, t) : S \text{ is a set of integers, } t \text{ is an integer, } \exists S' \subseteq S \text{ such that } \sum_{x \in S'} x = t\}.$$

The *partition problem* is defined as follows:

$$\text{PARTITION} = \{S : S \text{ is a set of integers, } \exists S' \subseteq S \text{ such that } \sum_{x \in S'} x = \sum_{y \in S \setminus S'} y\}.$$

- Prove that  $\text{SUBSETSUM} \leq_P \text{PARTITION}$ , i.e., in polynomial time, SUBSETSUM can be reduced to PARTITION.
- Prove that  $\text{PARTITION} \leq_P \text{SUBSETSUM}$ , i.e., in polynomial time, PARTITION can be reduced to SUBSETSUM.

**Question 5:** The *clique and independent set problem* is defined as follows:

$$\text{CLIQUEINDEPSET} = \{(G, K) : \text{graph } G \text{ contains a clique of size } K \text{ and } G \text{ contains an independent set of size } K\}.$$

Prove that  $\text{CLIQUE} \leq_P \text{CLIQUEINDEPSET}$ , i.e., in polynomial time, CLIQUE can be reduced to CLIQUEINDEPSET.

**Question 6:** Let  $\varphi$  be a Boolean formula in the variables  $x_1, x_2, \dots, x_n$ . We say that  $\varphi$  is in *conjunctive normal form* (CNF) if it is of the form

$$\varphi = C_1 \wedge C_2 \wedge \dots \wedge C_m,$$

where each  $C_i$ ,  $1 \leq i \leq m$ , is of the following form:

$$C_i = l_1^i \vee l_2^i \vee \dots \vee l_{k_i}^i.$$

Each  $l_j^i$  is a *literal*, which is either a variable or the negation of a variable.

The *satisfiability problem* is defined as follows:

$$\text{SAT} = \{\varphi : \varphi \text{ is in CNF-form and is satisfiable}\}.$$

Prove that  $\text{CLIQUE} \leq_P \text{SAT}$ , i.e., in polynomial time,  $\text{CLIQUE}$  can be reduced to  $\text{SAT}$ .