

# Testing the Quality of Manufactured Balls<sup>\*</sup>

Prosenjit Bose and Pat Morin

Carleton University, Ottawa, Canada, K1S 5B6  
{jit,morin}@scs.carleton.ca

**Abstract.** We consider the problem of testing the roundness of a manufactured ball, using the finger probing model of Cole and Yap [4]. When the center of the object is known, a procedure requiring  $O(n^2)$  probes and  $O(n^2)$  computation time is described. (Here  $n = |1/q|$ , where  $q$  is the quality of the object.) When the center of the object is not known, the procedure requires  $O(n^2)$  probes and  $O(n^4)$  computation time. We also give lower bounds that show that the number of probes used by these procedures is optimal.

## 1 Introduction

The field of metrology is concerned with measuring the quality of manufactured objects. A basic task in metrology is that of determining whether a given manufactured object is of acceptable quality. Usually this involves probing the surface of the object using a measuring device such as a coordinate measuring machine to get a set  $S$  of sample points, and then verifying, algorithmically, how well  $S$  approximates an ideal object.


A special case of this problem is determining whether an object is *round*, or *spherical*. For our purposes, an object  $I$  is *good* if there exists two concentric spheres  $I_{\text{in}}$  and  $I_{\text{out}}$  of radius  $1 - \epsilon$  and  $1 + \epsilon$ , respectively, such that  $I_{\text{in}}$  is entirely contained in  $I$  and  $I$  is entirely contained in  $I_{\text{out}}$ , and *bad* otherwise. We call the problem of deciding whether an object is good or bad the *roundness classification problem*. See Figure 1 for examples of good and bad objects.

In the field of computational geometry, the algorithmic side of the roundness classification problem has received considerable attention and efficient algorithms for testing the roundness of a set of 2D [1, 5, 6, 7, 9, 10, 12, 13, 14, 15] and 3D [6] sample points are known. However, very little research has been done on probing strategies for the roundness classification problem. Notable exceptions are the work by Mehlhorn, Shermer, and Yap [11], in which planar objects (disks) are considered, Bose and Morin [3], in which disks and cylinders are considered, and Fu and Yap [8] in which a probing strategy for finding the near-center of a  $d$ -dimensional ball using  $d(d + 1)$  probes is presented.

In this paper we describe strategies for testing the roundness of manufactured balls. (A ball is a solid object whose surface is a sphere.) We use the *finger probing*

---

<sup>\*</sup> This work was funded in part by the Natural Sciences and Engineering Research Council of Canada.



**Fig. 1.** Examples of (a) good and (b,c) bad objects.

model of Cole and Yap [4]. In this model, the measurement device can identify a point in the interior of  $I$  and can probe along any ray, i.e., determine the first point on the ray that intersects the boundary of  $I$ . The finger probing model is a reasonable abstract model of a coordinate measuring machine [16].

We describe a procedure for testing the roundness of a manufactured ball  $I$  using  $O(1/\text{qual}(I)^2)$  finger probes. Here  $|\text{qual}(I)|$  measures how far the object  $I$  is from the boundary between good and bad. When the center of  $I$  is known in advance, the procedure requires  $O(1/\text{qual}(I)^2)$  computation time. When a center of  $I$  is not known, the procedure requires  $O(1/\text{qual}(I)^4)$  computation time. As part of this procedure, we describe a technique for finding a near-center of a 3-dimensional ball that requires only 10 probes, thus providing an alternative to the procedure of Fur and Yap for the case  $d = 3$ . We also give a lower bound that shows our procedures are optimal, up to constant factors, in terms of the number of probes used.

The remainder of the paper is organized as follows: Section 2 gives definitions and notation used throughout the remainder of the paper. Section 3 describes a procedure for find a point near the center of an object. Section 4 discusses procedures for testing the roundness of an object. Section 5 gives a lower bound on the number of probes needed for this problem. Section 6 summarizes and suggests directions for future work.

## 2 Definitions, Notation, and Assumptions

In this section, we introduce definitions and notation used throughout the remainder of this paper, and state the assumptions we make on the object being tested. For the most part, notation and definitions are consistent with, or analogous to, [3, 11].

For a point  $p$ , we use the notation  $x(p)$ ,  $y(p)$ , and  $z(p)$  to denote the  $x$ ,  $y$ , and  $z$  coordinates of  $p$ , respectively. The letter  $O$  is used to denote the origin of the coordinate system. We use the notation  $\text{dist}(a, b)$  to denote Euclidean distance

between two objects. When  $a$  and  $b$  are not points,  $\text{dist}(a, b)$  is the minimum distance between all pairs of points in  $a$  and  $b$ . The angle formed by three points  $a$ ,  $b$ , and  $c$ , is denoted by  $\angle abc$ , and we always mean the smaller angle unless stated otherwise.

A *sphere* (*ball*) of radius  $r$  centered at a point  $c$  is the set of all points  $p$  such that  $\text{dist}(p, c) = r$  ( $\text{dist}(p, c) \leq r$ ). A sphere (ball) of with radius  $r = 1$  is called a *unit sphere* (*unit ball*). Two spheres or balls are said to be *concentric* if they are centered at the same point.

An *object*  $I$  is defined to be any compact simply connected subset of 3-space, with boundary denoted by  $\text{bd}(I)$ . For a point  $p$ , we use  $R(p, I)$  and  $r(p, I)$  to denote the maximal and minimal distance, respectively, from  $p$  to a point in  $\text{bd}(I)$ . I.e.,

$$R(p, I) = \max\{\text{dist}(p, p') : p' \in \text{bd}(I)\} \quad (1)$$

$$r(p, I) = \min\{\text{dist}(p, p') : p' \in \text{bd}(I)\} . \quad (2)$$

For a point  $p$ , let

$$\text{qual}(p, I) = \min\{r(p, I) - (1 - \epsilon), (1 + \epsilon) - R(p, I)\} \quad (3)$$

and let

$$\text{qual}(I) = \max_{p \in I} \text{qual}(p, I) . \quad (4)$$

Any point  $c_I$  with  $\text{qual}(c_I, I) = \text{qual}(I)$  is called a *center* of  $I$ . Note that there may be more than one point with this property, i.e., the center of  $I$  is not necessarily unique.

The value  $\text{qual}(I)$  is called the *quality* of the object  $I$ , since it measure the maximum deviation of  $I$  from a ball of unit radius. An object  $I$  with  $\text{qual}(I) > 0$  is *good* while an object  $I$  with  $\text{qual}(I) < 0$  is *bad*. A procedure that determines whether an object is good or bad is called a *roundness classification procedure*.

In order to have a roundness classification procedure that is correct and that terminates, it is necessary to make some assumptions about the object  $I$  being tested. The following assumption is referred to as the *minimum quality assumption*, and refers to the fact that the manufacturing process can guarantee that manufactured objects have a minimum quality (although perhaps not enough to satisfy our roundness criteria). The constant  $1/30$  in the assumption is easily met by current manufacturing processes.

**Assumption 1** There exists two concentric balls,  $I_{\text{in}}$  and  $I_{\text{out}}$ , with radii  $1 - \delta$  and  $1 + \delta$ , respectively, such that  $I_{\text{in}} \subseteq I \subseteq I_{\text{out}}$ , for some  $\delta < 1/30$ .

The minimum quality assumption alone is not sufficient. If the object under consideration contains oddly shaped recesses, then it may be the case that these recesses can not be probed using finger probes. We say that an on object  $I$  is *star-shaped* if there exists a point  $k \in I$  such that for any point  $p \in I$ , the line segment joining  $k$  and  $p$  is a subset of  $I$ . We call the set of all points with this property the *kernel* of  $I$ . There is a region about the center  $c_I$  of  $I$  that is of

particular interest. The following assumption ensures that all points in  $\text{bd}(I)$  can be probed by directing probes at a point close to  $c_I$ .

**Assumption 2** Let  $c_I$  be any center of  $I$ .  $I$  is a star-shaped object, and its kernel contains all points  $p$  such that  $\text{dist}(c_I, p) \leq \alpha$ , for some constant  $1 - \delta > \alpha > 2\delta$ .

### 3 Finding a Near-Center

In this section we describe a procedure for finding a point close to the center,  $c_I$ , of  $I$ . A *near-center* of  $I$  is any point  $c_0$  such that  $\text{dist}(c_0, c_I) \leq 2\delta$ . Our procedure uses three simple subroutines  $X(p)$ ,  $Y(p)$  and  $Z(p)$ . These subroutines perform two probes directed at  $p$ . The two probes come from opposite directions, and are parallel to the  $x$ ,  $y$ , and  $z$  axes, respectively. If the two probes contact  $I$  at points  $a$  and  $b$ , then the subroutines return  $(a + b)/2$ , i.e., the midpoint between  $a$  and  $b$ . If the probes do not contact  $I$  then the routines return the point  $p$ . Pseudocode is given in Procedure 1.

---

**Procedure 1** Returns a near-center given a point  $p_0 \in I$ .

---

```

1:  $p_1 \leftarrow X(p_0)$ 
2:  $p_2 \leftarrow Y(p_1)$ 
3:  $p_3 \leftarrow Z(p_2)$ 
4:  $p_4 \leftarrow X(p_3)$ 
5:  $p_5 \leftarrow Y(p_4)$ 
6: return  $p_5$ 

```

---

**Theorem 1** Let  $I$  be an object with center  $c_I$  and satisfying Assumption 1. Then 10 probes and constant computation time suffice to find a point  $c_0$  such that  $\text{dist}(c_I, c_0) \leq 2\delta$ .


*Proof.* The proof involves using Assumption 1 to bound the values of the  $x$ ,  $y$  and  $z$  coordinates of  $p_{1-5}$  and is rather long. It is omitted due to space constraints but can be found in the full version of the paper [2].  $\square$

### 4 Testing Quality

Once a center or near-center of  $I$  is known, we can obtain an approximation of the surface of  $I$  by directing probes at this (near) center. In this section, we first describe a strategy for directing probes at the (near) center. We then describe the entire quality testing procedure for the case when the center of  $I$  is known in advance. Finally, we describe the procedure for the case when the center of  $I$  is not known in advance. Proving the correctness of our procedures involves bounding the maximum error in our approximation of the surface.

## 4.1 The Probing Strategy

In this section, we describe a probing strategy for taking  $\Theta(n^2)$  probes directed at a point  $p$ , where  $n$  is an even positive integer. The strategy is designed so that for any direction  $d$ , there is a probe in some direction “not too far” from  $d$ . Refer to Figure 2 for an illustration of what follows.



**Fig. 2.** Illustration of (a) spherical coordinates and (b) partitioning the sphere into slices and pieces.

Consider the spherical coordinates  $(\phi, \rho)$  of the unit sphere centered at  $p$ , where angles  $\phi$  and  $\rho$  are in the set  $[0, 2\pi)$ . We first divide the sphere into  $n$  parallel slices,  $s_0, \dots, s_{n-1}$ , such that slice  $s_i$  contains all points where  $\rho \in [i\pi/n, (i+1)\pi/n]$ . Each slice  $s_i$  is further subdivided into  $m_i = \lceil 2n \max\{\sin(i\pi/n), \sin((i+1)\pi/n)\} \rceil$  similar pieces,  $s_{i0}, \dots, s_{i(m_i-1)}$ , such that piece  $s_{ij}$  contains all points in  $s_i$  where  $\phi \in [j\pi/m_i, (j+1)\pi/m_i]$ . We define the center of a piece  $s_{ij}$  as the point with spherical coordinates  $((2i+1)\pi/2n, (2j+1)\pi/2m_i)$ .

**Lemma 1** *Let  $a$  be any point in  $s_{ij}$ , and let  $b$  be the center of  $s_{ij}$ . Then  $\angle apb \leq \pi/n$ .*

*Proof.* The proof is straightforward. It is omitted due to space constraints but can be found in the full version of the paper [2].  $\square$

**Lemma 2**  $\sum_{i=0}^{n-1} m_i \in \Theta(n^2)$ , i.e., the partitioning of the sphere described above contains  $\Theta(n^2)$  pieces.

*Proof.* That the number of pieces is  $O(n^2)$  follows from the inequality  $\sin(\tau) \leq 1$ . That the number of pieces is  $\Omega(n^2)$  follows from the inequality  $\sin(\tau) \geq 2\tau/\pi$ , for  $\tau \in [0, \pi/2]$ .  $\square$

For some center or near center  $c$ , our probing strategy involves directing probes along each of the half lines with an endpoint at  $c$  and passing through

the center of each piece of the sphere centered at  $c$ . In the remainder of the paper, we will use the notation  $\text{probe}(n, c)$  to denote the set of probes obtained when using this strategy.

## 4.2 The Simplified Procedure

In this section we describe a simplified roundness classification procedure that assumes that we know the object being tested is centered at the origin,  $O$ . Our roundness classification procedure (Procedure 2) tests the roundness of an object  $I$  by taking a set  $S$  of probes in the manner described in the previous section. The procedure repeatedly doubles the number of probes until either (1) a set of sample points is found that proves that  $I$  is a bad object, in which case  $I$  is rejected, or (2) the quality of the set of sample points is “significantly larger” than 0, in which case we can prove that  $\text{qual}(I) > 0$ .

---

**Procedure 2** Tests the roundness of the object  $I$  centered at the origin.

---

```

1:  $r \leftarrow 1$ 
2:  $R \leftarrow 1$ 
3:  $n \leftarrow n_0$ 
4:  $\Delta \leftarrow f(n) \in O(1/n)$ 
5: repeat
6:    $S \leftarrow \text{probe}(n, O)$ 
7:   if  $\exists p \in S : \text{dist}(p, O) > 1 + \epsilon$  or  $\text{dist}(p, O) < 1 - \epsilon$  then
8:     return REJECT
9:   end if
10:   $r \leftarrow 1 - \epsilon + \Delta$ 
11:   $R \leftarrow 1 + \epsilon - \Delta$ 
12:   $n \leftarrow 2n$ 
13:   $\Delta \leftarrow f(n)$ 
14: until  $\forall p \in S : \text{dist}(p, O) < R$  and  $\text{dist}(p, O) > r$ 
15: return ACCEPT

```

---

The function  $f(n)$  that appears in the procedure is defined as

$$f(n) = \frac{1}{n} \left( \frac{1 + \delta}{\alpha} \right) (\alpha^2 + 1 + 2\delta + \delta^2)^{\frac{1}{2}} ,$$

and the constant  $n_0$  is defined as

$$n_0 = \lceil \pi / \arctan(\alpha / (1 + \delta)) \rceil .$$

**Lemma 3** *Let  $I$  be an object with center  $c_I = O$ . Let  $S = \text{probe}(n, c_I)$ , for any  $n \geq n_0$ . Then for any point  $p \in \text{bd}(I)$ , there exists a point  $p' \in S$  such that  $\text{dist}(p, p') \leq f(n)$ .*


*Proof.* We will bound  $|x(p) - x(p')|$ ,  $|y(p) - y(p')|$ , and  $|z(p) - z(p')|$ . Refer to Figure 3 for an illustration of what follows.

By Lemma 1, there exists a point  $p' \in S$  such that

$$\angle pc_I p' \leq \pi/n . \quad (5)$$

By orienting the coordinate system so that the plane  $z = 0$  passes through  $p$ ,  $p'$  and  $c_I$ , we can assume, wlog, that

$$|z(p) - z(p')| = 0 . \quad (6)$$



**Fig. 3.** Constraints on the position of  $p'$ . The point  $p'$  must be in the shaded region, and  $\text{dist}(p, p')$  is maximized when  $p'$  is placed as shown.

Next note that we can rotate the coordinate system about the  $z$ -axis so that we can assume, wlog, that  $x(p) = 0$ , and  $1 - \delta \leq y(p) \leq 1 + \delta$ . Assumption 1 ensures that  $\text{dist}(O, p') \leq 1 + \delta$ . So, by (5), an upper bound on  $|x(p) - x(p')|$  is

$$|x(p) - x(p')| = |x(p')| \quad (7)$$

$$\leq (1 + \delta) \sin(\pi/n) \quad (8)$$

$$\leq (1 + \delta)\pi/n . \quad (9)$$

Since  $\angle pc_I p' \leq \pi/n$ , the point  $p'$  must lie in the cone defined by the inequality

$$y(p') \geq |x(p')| (\cos(\pi/n) / \sin(\pi/n)) . \quad (10)$$

Next we note that the slope of the line through  $p'$  and  $p$  must be in the range  $[-y(p)/\alpha, y(p)/\alpha]$ , otherwise Assumption 2 is violated. If  $n \geq n_0 = \pi / \arctan(\alpha/y(p))$ , the region in which  $p'$  can be placed is bounded, and  $|y(p) - y(p')|$  is maximized when  $p'$  lies on one of the bounding lines

$$f_l(x) = xy(p)/\alpha + y(p)$$

$$f_r(x) = -xy(p)/\alpha + y(p)$$

Since both lines are symmetric about  $x = 0$  we can assume that  $x(p')$  lies on  $f_l$ , giving us

$$\begin{aligned}
|y(p) - y(p')| &\leq |y(p) - f_l(x(p'))| \\
&= |x(p')y(p)/\alpha| \\
&\leq |x(p')(1 + \delta)/\alpha| \\
&= (1 + \delta)^2\pi/\alpha n
\end{aligned} \tag{11}$$

Plugging (9), (11), and (6) into the Euclidean distance formula and simplifying yields the desired result.  $\square$

**Theorem 2** *There exists a roundness classification procedure that can correctly classify any object  $I$  with center  $c_I = O$  and satisfying Assumptions 1 and 2 using  $O(1/\text{qual}(I)^2)$  probes and  $O(1/\text{qual}(I)^2)$  computation time.*

*Proof.* We begin by showing that the Procedure 2 is correct. We need to show that the procedure never rejects a good object and never accepts a bad object. The former follows from the fact that the procedure only ever rejects an object when it finds a point on the object's boundary whose distance is less than  $1 - \epsilon$  or greater than  $1 + \epsilon$  from  $c_I$ . To show the latter, we note that Lemma 3 implies that there is no point in  $\text{bd}(I)$  that is of distance greater than  $f(n)$  from all points in  $S$ . The procedure only accepts  $I$  when the distance of all points in  $S$  from  $c_I$  are in the range  $[1 - \epsilon + f(n), 1 + \delta - f(n)]$ . Therefore, if the procedure accepts  $I$ , the distance of all points in  $\text{bd}(I)$  from  $c_I$  is in the range  $[1 - \epsilon, 1 + \delta]$ , i.e., the object is good.

Next we prove that the running time is  $O(1/\text{qual}(I)^2)$ . First we observe that  $f(n) \in O(1/n)$ . Next, note that the computation time and number of probes used during each iteration is linear with respect to the value of  $n$ , and the value of  $n$  doubles after each iteration. Thus, asymptotically, the computation time and number of probes used are dominated by the value of  $n^2$  during the last iteration. There are two cases to consider.

**Case 1:** Procedure 2 accepts  $I$ . In this case, the procedure will certainly terminate once  $\Delta \leq \text{qual}(I)$ . This takes  $O(\log(1/\text{qual}(I)))$  iterations. During the final iteration,  $n \in O(1/\text{qual}(I))$ .

**Case 2:** Procedure 2 rejects  $I$ . In this case, there is a point on  $\text{bd}(I)$  at distance  $\text{qual}(I)$  outside the circle with radius  $1 + \epsilon$  centered at  $O$ , or there is a point in  $\text{bd}(I)$  at distance  $\text{qual}(I)$  inside of the circle with radius  $1 - \epsilon$  centered at  $O$ . In either case, Lemma 3 ensures that the procedure will find a bad point within  $O(\log |1/\text{qual}(I)|)$  iterations. During the final iteration,  $n \in O(|1/\text{qual}(I)|)$ .  $\square$

### 4.3 The Full Procedure

In the more general (and realistic) version of the roundness classification problem, we do not know the center of the object being tested. However, Theorem 1 allows us to use this procedure anyhow. The significance of Theorem 1 is that it allows us to find a near-center,  $c_0$ , of  $I$ . As the following lemma shows, knowing



a near-center is almost as useful as knowing the true center. Before we state the lemma, we need the following definitions.

$$f'(n) = \frac{1}{n} \left( (1 + 3\delta)^2 \pi^2 + \frac{(1 + 3\delta)^4 \pi^2}{(\alpha - 2\delta)^2} \right)^{\frac{1}{2}}$$

$$n'_0 = \lceil \pi / \arctan(\alpha / (1 + 3\delta)) \rceil$$

**Lemma 4** *Let  $I$  be an object with center  $c_I$  and near-center  $c_0$ , and satisfying Assumptions 1 and 2. Let  $S = \text{probe}(n, c_0)$ , for any  $n \geq n'_0$ , where  $\text{dist}(c_0, c_I) \leq 2\delta$ . Then for any point  $p \in \text{bd}(I)$ , there exists a point  $p' \in S$  such that  $\text{dist}(p, p') \leq f'(n)$ .*

*Proof.* The proof is almost a verbatim translation of the proof of Lemma 3, except that we assume, wlog, that  $c_0 = O$ . With this assumption we derive the bounds

$$|x(p) - x(p')| \leq (1 + 3\delta)(\pi/n)$$

$$|y(p) - y(p')| \leq (1 + 3\delta)^2 \pi / n (\alpha - 2\delta)$$

Substituting these values into the formula for the Euclidean distance and simplifying yields the desired result.  $\square$

**Lemma 5** *Let  $I$  be an object with center  $c_I$  and near-center  $c_0$  and satisfying Assumptions 1 and 2. Let  $S = \text{probe}(n, c_0)$ , for any  $n \geq n'_0$ , and let  $c_S$  be a center of  $S$ . Then*

$$R(c_S, S) \leq R(c_S, I) \leq R(c_S, S) + f'(n) ,$$

$$r(c_S, S) - f'(n) \leq r(c_S, I) \leq r(c_S, S) .$$

*Proof.* The lemma follows from (1), (2) and Lemma 4. The proof is omitted due to space constraints but can be found in the full version of the paper [2].  $\square$

**Lemma 6** *Let  $I$  be an object with center  $c_I$  and near-center  $c_0$  and satisfying Assumptions 1 and 2. Let  $S = \text{probe}(n, c_0)$ , for any  $n \geq n'_0$ . Then  $\text{qual}(S) - f'(n) \leq \text{qual}(I) \leq \text{qual}(S)$*

*Proof.* The lemma follows from (4) and Lemma 5. The proof is omitted due to space constraints but can be found in the full version of the paper [2].  $\square$

In [6], an algorithm is described that determines, given a set  $S$  of points in 3-space, the minimum value of  $\epsilon$  such that there exists two concentric closed balls  $I_{\text{in}}$  and  $I_{\text{out}}$ , with radii  $1 - \epsilon$  and  $1 + \epsilon$ , respectively, such that  $S \cap I_{\text{in}} = \emptyset$  and  $S \cap I_{\text{out}} = S$ . The running time of the algorithm is  $O(|S|^2)$ . Combining Lemma 4 with this algorithm, we obtain the following result.

**Theorem 3** *There exists a roundness classification procedure that can correctly classify any object  $I$  satisfying Assumptions 1 and 2 using  $O(1/\text{qual}(I)^2)$  probes and  $O(1/\text{qual}(I)^4)$  computation time.*

*Proof.* We make the following modifications to Procedure 2. In Line 3, we set the value of  $n$  to  $n'_0$ . In Lines 4 and 13, we replace  $f(n)$  with  $f'(n)$ . In Line 6 we directed our probes at  $c_0$  rather than  $O$ . In Lines 7 and 14, we replace the simple test with a call to the algorithm of [6].

Lemma 6 ensures that the procedure never accepts a bad object and never rejects a good object. i.e., the procedure is correct. The procedure terminates once  $f'(n) < |\text{qual}(I)|$ . This happens after  $O(\log |1/\text{qual}(I)|)$  iterations, at which point  $n \in O(|1/\text{qual}(I)|)$ .  $\square$

## 5 Lower Bounds

In this section, we give a lower bound that shows that any correct roundness classification procedure for spheres requires, in the worst case,  $\Omega(1/\text{qual}(I)^2)$  probes to determine if  $I$  is good or bad. The lower bound uses an adversary argument to show that if a procedure uses  $o(1/\text{qual}(I)^2)$  probes, then an adversary can orient a bad object so that its defects are “hidden” from all the probes, making the bad object indistinguishable from a similar good object.

**Lemma 7** *Let  $S$  be a set of  $n^2$  points on the unit sphere. Then, there exists a spherical cap  $c$  with radius  $1/n$  such that  $c$  contains no points of  $S$*

*Proof.* Consider the convex hull of  $S$ , which has at most  $2n^2 - 4$  faces. The plane passing through a face defines a spherical cap, and the union of these caps cover the entire sphere, a surface area of  $4\pi$ . By the pigeonhole principle, some face  $f$  must define a cap  $c$  with surface area at least  $2\pi/n^2$ . Furthermore, since  $f$  is part of the convex hull, there are no other points of  $S$  in this cap. The surface area of  $c$  obeys the inequality  $\text{sa}(c) \leq 2\pi r^2$ , where  $r$  is the radius of  $c$ . Thus, we have the inequalities

$$2\pi r^2 \geq \text{sa}(c) \geq 2\pi/n^2 ,$$


yielding  $r \geq 1/n$ .  $\square$

**Theorem 4** *Any roundness classification procedure that is always correct requires, in the worst case,  $\Omega(|1/\text{qual}(I)|^2)$  probes to classify a object  $I$  with center  $c_I = O$  and satisfying Assumptions 1 and 2.*

*Proof.* We prove the theorem by exhibiting two objects  $I$  and  $I'$  with  $\text{qual}(I) = \psi = -\text{qual}(I')$ , for any  $0 \leq \psi \leq \epsilon$ , such that  $I$  and  $I'$  cannot be distinguished by any algorithm that uses  $o(|1/\text{qual}(I)|)$  probes.

The object  $I$  is a perfect circle with radius  $1 - \epsilon + \psi$ . The object  $I'$  is similar to  $I$ , except that it contains a conic recess of depth  $4\psi$  that removes a circle of diameter  $\alpha 8\psi$  from the surface of  $I$  (see Figure 4). Note that  $\text{qual}(I) = \psi$  and  $\text{qual}(I') = -\psi$ , and that for  $\alpha = 1/9$ ,  $\delta \leq 1/21$ , and  $\psi \leq \epsilon \leq \delta$ ,  $I$  and  $I'$  satisfy Assumptions 1 and 2.

Assume by way of contradiction that there exists a roundness classification procedure  $\mathcal{P}$  that always accepts  $I$  and always rejects  $I'$  using  $o(1/\psi^2)$  probes.



**Fig. 4.** An example of two objects,  $I$  and  $I'$  that cannot be distinguished using  $o(1/\text{qual}(I)^2)$  probes.

Let  $S$  be the set of probes made by  $\mathcal{P}$  in classifying  $I$ . By Lemma 7, there exists a spherical cap  $c$  on the surface of  $I$  with radius  $\omega(\psi)$  such that  $c$  contains no point of  $S$ .<sup>1</sup> Therefore, for sufficiently small  $\psi$ ,  $c$  has diameter larger than  $\alpha 8\psi$  and an adversary can orient  $I'$  so that  $\mathcal{P}$  does not direct any probes at the conic recess in  $I'$ . The results of probes performed by, and therefore the actions of  $\mathcal{P}$ , would be the same for  $I$  and  $I'$ . But this is a contradiction, since we assumed that  $\mathcal{P}$  always correctly classifies both  $I$  and  $I'$ .  $\square$

## 6 Conclusions

We have described the first roundness classification procedure for balls and given lower bounds that show that the procedure is optimal in terms of the number of probes used. In the case when the center of the object is known in advance, the procedure is also optimal in terms of computation time.

When the center of the object is not known, our procedure would benefit significantly from an improved algorithm for testing the roundness of a 3-dimensional point set. The algorithm in [6], which we rely on, solves the problem by first constructing the Voronoi diagram of the point set, which can have quadratic complexity in the worst case. A subquadratic time algorithm is still an important open problem.

## References

- [1] P. Agarwal, B. Aronov, and M. Sharir. Line transversals of balls and smallest enclosing cylinder in three dimensions. In *8th ACM-SIAM Symposium on Data Structures and Algorithms (SODA)*, pages 483–492, 1997.

<sup>1</sup> This is, unfortunately, a confusing use of asymptotic notation. The reader should keep in mind that it is the behaviour of the radius as  $\psi$  goes to zero that is being considered.

- [2] P. Bose and P. Morin. Testing the quality of manufactured balls. Technical Report TR-98-08, Carleton University School of Computer Science, 1998.
- [3] P. Bose and P. Morin. Testing the quality of manufactured disks and cylinders. In *Proceedings of the Ninth Annual International Symposium on Algorithms and Computation (ISAAC'98)*, pages 129–138, 1998.
- [4] R. Cole and C. K. Yap. Shape from probing. *Journal of Algorithms*, 8:19–38, 1987.
- [5] M. deBerg, P. Bose, D. Bremner, S. Ramaswami, and G. Wilfong. Computing constrained minimum-width annuli of point sets. In *Proceedings of the 5th Workshop on Data Structures and Algorithms*, pages 25–36, 1997.
- [6] C. A. Duncan, M. T. Goodrich, and E. A. Ramos. Efficient approximation and optimization algorithms for computational metrology. In *8th ACM-SIAM Symposium on Data Structures and Algorithms (SODA)*, pages 121–130, 1997.
- [7] H. Ebara, N. Fukuyama, H. Nakano, and Y. Nakanishi. Roundness algorithms using the Voronoi diagrams. In *1st Canadian Conference on Computational Geometry*, page 41, 1989.
- [8] Q. Fu and C. K. Yap. Computing near-centers in any dimension. Unpublished manuscript, 1998.
- [9] J. Garcia and P. A. Ramos. Fitting a set of points by a circle. In *ACM Symposium on Computational Geometry*, 1997.
- [10] V. B. Le and D. T. Lee. Out-of-roundness problem revisited. *IEEE Transactions on Pattern Analysis and Machine Intelligence*, 13(3):217–223, 1991.
- [11] K. Mehlhorn, T. Shermer, and C. Yap. A complete roundness classification procedure. In *ACM Symposium on Computational Geometry*, pages 129–138, 1997.
- [12] P. Ramos. Computing roundness in practice. In *European Conference on Computational Geometry*, pages 125–126, 1997.
- [13] U. Roy and X. Zhang. Establishment of a pair of concentric circles with the minimum radial separation for assessing rounding error. *Computer Aided Design*, 24(3):161–168, 1992.
- [14] E. Schomer, J. Sellen, M. Teichmann, and C. K. Yap. Efficient algorithms for the smallest enclosing cylinder. In *8th Canadian Conference on Computational Geometry*, pages 264–269, 1996.
- [15] K. Swanson. An optimal algorithm for roundness determination on convex polygons. In *Proceedings of the 3rd Workshop on Data Structures and Algorithms*, pages 601–609, 1993.
- [16] C. K. Yap. Exact computational geometry and tolerancing metrology. In David Avis and Jit Bose, editors, *Snapshots of Computational and Discrete Geometry, Vol. 3*. 1994.